



Lie group analysis for short pulse equation

Mehdi Nadjafikhah^{*a}

^aDepartment of Pure Mathematics, School of Mathematics, Iran University of Science and Technology, Narmak, Tehran, 16846-13114, Iran.

ABSTRACT: In this paper, the classical Lie symmetry analysis and the generalized form of Lie symmetry method are performed for a general short pulse equation. The point, contact and local symmetries for this equation are given. In this paper, we generalize the results of H. Liu and J. Li [2], and add some further facts, such as an optimal system of Lie symmetry subalgebras and two local symmetries.

Review History:

Received:12 May 2020

Revised:26 June 2020

Accepted:27 June 2020

Available Online:01 September 2020

Keywords:

Lie symmetry analysis
General short pulse equation
Invariant solution
Local symmetry

AMS Subject classifications:

17B80, 22E70, 35C05

1. Introduction

Non-linear PDEs arising in many applied fields like Biology, Fluid Mechanics, Plasma Physics and Optics, systems of impulse and Neural Networks, etc. and exhibit a rich variety of non-linear phenomena. The investigation of the exact solutions plays an important role in the study of non-linear systems. In this paper, we find Lie point symmetries, third order local symmetries, an optimal system of these two type symmetries, and corresponding invariant solutions for a general short pulse equation:

$$\text{SPE} : u_{xt} = \alpha u + \frac{\beta}{3}(u^3)_{xx}, \quad (1)$$

where $u = u(x, t)$ is the unknown real function and subscripts denote differentiation w.r.t. x and t ; α and β are non-zero real parameters.

This general SPE was derived by T. Schafer and C.E. Wayne [8, p.94] as a model equation describing the propagation of ultra-short light pulses in silica optical fibres. In [8, 7], many results are obtained about the special SPE:

$$u_{xt} = u + \frac{1}{6}(u^3)_{xx}. \quad (2)$$

^{*}Corresponding author.

E-mail addresses: m_nadjafikhah@iust.ac.ir

2. Lie contact and point symmetries

Let $J^1 = J^1(\mathbb{R}^2, \mathbb{R})$ be the jet space with coordinates (x, t, u, u_x, u_t) . Let

$$\mathbf{v} = \xi \partial_x + \tau \partial_t + \eta \partial_u + \eta^x \partial_{u_x} + \eta^t \partial_{u_t}, \tag{3}$$

be an infinitesimal Lie contact symmetry of (1), where ξ, τ and η are smooth functions $J^1(\mathbb{R}^2, \mathbb{R}) \rightarrow \mathbb{R}$, and

$$Q = \xi u_x + \tau u_t - \eta, \tag{4}$$

be the characteristic function of (4). Thus

$$\begin{aligned} \xi &= Q_{u_x}, & \tau &= Q_{u_t}, & \eta &= u_x Q_{u_x} + u_t Q_{u_t} - Q, \\ \eta^x &= -Q_x - u_x Q_u, & \eta^t &= -Q_t - u_t Q_u. \end{aligned} \tag{5}$$

Then, the (3) is an infinitesimal Lie contact symmetry of the (1) if and only if

$$\mathbf{v}^{(2)} \left(u_{xt} - \alpha u - \frac{1}{3} \beta (u^3)_{xx} \right) = 0, \quad u_{xt} = \alpha u + \frac{1}{3} \beta (u^3)_{xx}, \tag{6}$$

where $\mathbf{v}^{(2)}$ is the second prolongation of \mathbf{v} :

$$\mathbf{v}^{(2)} = Q \partial_u + D_x Q \partial_{u_x} + D_t Q \partial_{u_t} + D_x^2 Q \partial_{u_{xx}} + D_x D_t Q \partial_{u_{xt}} + D_t^2 Q \partial_{u_{tt}}, \tag{7}$$

where D_x and D_t are total derivative w.r.t x and t , respectively.

By substituting $u_{x,t}$ from second equation of (7) in the first equation, we find a polynomial of u_{xx} and u_{tt} with functional coefficients of (x, t, u, u_x, u_t) . Its coefficients must be zero:

$$\begin{aligned} Q_{u_x, u_x} &= 0, & Q_{u_x, u_t} &= 0, & Q_{u_t, u_t} &= 0, \\ u_t Q_{u, u_x} + \alpha u Q_{u_x, u_x} + Q_{t, u_x} &= 0, & \alpha u Q_{u_t, u_t} + u_x Q_{u, u_t} + Q_{x, u_t} &= 0, \\ u_x^2 Q_{u, u} + 2u_x Q_{x, u} + Q_{xx} + \alpha u (\alpha Q_{u_t, u_t} + 2u_x Q_{u, u_t} + 2Q_{x, u_t}) &= 0, \\ u_t Q_{u, u_t} - 5u_x Q_{u, u_x} - 5Q_{x, u_x} + Q_{t, u_t} - 4u Q_{u_x, u_t} - 2u_x Q_u &= 0, \\ u_x u_t Q_{u, u} + u_x Q_{t, u} + u_t Q_{x, u} + Q_{x, t} \\ + \alpha u (u_t Q_{u, u_t} + Q_{t, u_t} + Q_{x, u_x} + Q_u) + \alpha (u_x Q_{u_x} + u_t Q_{u_t} - Q) &= 0. \end{aligned} \tag{8}$$

After solving the determining system (8), one finds that

$$Q = c_1 u_x + c_2 u_t + c_3 (x u_x - t u_t - 3u); \tag{9}$$

where, c_1, c_2 and c_3 are arbitrary constants. Therefore,

Theorem 1. *The SPE (1) has a 3-dimensional Lie algebra \mathfrak{g} of point symmetries, generated by the infinitesimal generators*

$$\mathbf{v}_1 = \partial_x, \quad \mathbf{v}_2 = \partial_t, \quad \mathbf{v}_3 = x \partial_x - t \partial_t + 3u \partial_u, \tag{10}$$

and commuting table

$[\cdot, \cdot]$	\mathbf{v}_1	\mathbf{v}_2	\mathbf{v}_3	(11)
\mathbf{v}_1	0	0	\mathbf{v}_1	
\mathbf{v}_2	0	0	$-\mathbf{v}_2$	
\mathbf{v}_3	$-\mathbf{v}_1$	\mathbf{v}_2	0	

The SPE (1) has not any non-point contact symmetry.

3. Invariant solutions and its classification

The one-parameter groups G_i generated by the base of \mathfrak{g} are as follows:

$$\begin{aligned} G_1 &: \exp(\varepsilon \mathbf{v}_1) \cdot (x, t, u) = (x + \varepsilon, t, u), \\ G_2 &: \exp(\varepsilon \mathbf{v}_2) \cdot (x, t, u) = (x, t + \varepsilon, u), \\ G_3 &: \exp(\varepsilon \mathbf{v}_3) \cdot (x, t, u) = (e^\varepsilon x, e^{-\varepsilon} t, e^{3\varepsilon} u), \end{aligned} \tag{12}$$

where ε is a real number.

Since each group G_i is a symmetry group of SPE (1) and if $u = f(x, y)$ is a solution of the SPE (1), so are the following functions

$$u = f(x + \varepsilon, t), \quad u = f(x, t + \varepsilon), \quad u = f(e^\varepsilon x, e^{-\varepsilon} t, e^{-3\varepsilon} u), \tag{13}$$

where ε is an arbitrary real number. Thus, for the arbitrary combination $\mathbf{v} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 \in \mathfrak{g}$, the SPE (1) has the following solution:

$$u = f(e^{\varepsilon_3} x + \varepsilon_1, e^{-\varepsilon_3} t + \varepsilon_2, e^{-3\varepsilon_3} u), \tag{14}$$

where ε_i are arbitrary real numbers.

Let G be the symmetry Lie group of SPE (1). Now G operates on the set of solutions S of SPE (1), and $s \cdot G$ be the orbit of s , and H be a subgroup of G . Invariant H -solutions $s \in S$ are characterized by equality $s \cdot S = \{s\}$. If $h \in G$ is a transformation and $s \in S$, then

$$h \cdot (s \cdot H) = (h \cdot s) \cdot (hHh^{-1}). \tag{15}$$

Consequently, every invariant H -solution s transforms into an invariant hHh^{-1} -solution (Proposition 3.6 of [5]). Therefore, different invariant solutions are found from similar subgroups of G . Thus, the classification of invariant H -solutions is reduced to the problem of the classification of subgroups of G , up to similarity. An optimal system of s -dimensional subgroups of G is a list of conjugacy inequivalent s -dimensional subgroups of G with the property that any other subgroup is conjugate to precisely one subgroup in the list. Similarly, a list of s -dimensional sub-algebras forms an optimal system if every s -dimensional sub-algebra of \mathfrak{g} is equivalent to a unique member of the list under some element of the adjoint representation: $\tilde{\mathbf{h}} = \text{Ad}(g) \cdot \mathbf{h}$. Let \tilde{H} and \tilde{H} be connected, s -dimensional Lie subgroups of the Lie group G with corresponding Lie sub-algebras \mathbf{h} and $\tilde{\mathbf{h}}$ of the Lie algebra \mathfrak{g} . Then $\tilde{H} = gHg^{-1}$ are conjugate subgroups if and only if $\tilde{\mathbf{h}} = \text{Ad}(g) \cdot \mathbf{h}$ are conjugate sub-algebras (Proposition 3.7 of [5]). Thus, the problem of finding an optimal system of subgroups is equivalent to that of finding an optimal system of sub-algebras, and so we concentrate on it.

For the one-dimensional sub-algebras, the classification problem is essentially the same as the problem of classifying the orbits of the adjoint representation, since each one-dimensional sub-algebra is determined by a nonzero vector in Lie algebra symmetries of SPE (1) and so to "simplify" it as much as possible. The adjoint action is given by the Lie series

$$\text{Ad}(\exp(\varepsilon\mathbf{v}_i)\mathbf{v}_j) = \mathbf{v}_j - \varepsilon[\mathbf{v}_i, \mathbf{v}_j] + \frac{\varepsilon^2}{2}[\mathbf{v}_i, [\mathbf{v}_i, \mathbf{v}_j]] - \dots, \tag{16}$$

where $i, j = 1, \dots, 3$. Let $F_i^\varepsilon : \mathfrak{g} \rightarrow \mathfrak{g}$ defined by $\mathbf{v} \mapsto \text{Ad}(\exp(\varepsilon\mathbf{v}_i)\mathbf{v})$, for $i = 1, \dots, 3$. Therefore, if $\mathbf{v} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3 \in \mathfrak{g}$, then

$$\begin{aligned} F_i^{\varepsilon_1}(\mathbf{v}) &= (c_1 + \varepsilon_1 c_3)\mathbf{v}_1 + c_2\mathbf{v}_2 + c_3\mathbf{v}_3, \\ F_i^{\varepsilon_2}(\mathbf{v}) &= c_1\mathbf{v}_1 + (c_2 + \varepsilon_2 c_3)\mathbf{v}_2 + c_3\mathbf{v}_3, \\ F_i^{\varepsilon_3}(\mathbf{v}) &= e^{-\varepsilon_3} c_1\mathbf{v}_1 + e^{\varepsilon_3} c_2\mathbf{v}_2 + c_3\mathbf{v}_3. \end{aligned} \tag{17}$$

Applying these transformations, one can show that

Theorem 2. *A one-dimensional optimal system of \mathfrak{g} is*

$$\mathbf{v}_1 + a\mathbf{v}_2, \quad b\mathbf{v}_1 + \mathbf{v}_2, \quad \mathbf{v}_3, \tag{18}$$

where a and b are real constants; and, a two-dimensional optimal system of \mathfrak{g} is given by

$$\mathbf{v}_1, \mathbf{v}_2, \quad \mathbf{v}_1, \mathbf{v}_3, \quad \mathbf{v}_2, \mathbf{v}_3. \tag{19}$$

4. Local symmetries of SPE

One can generalize one-parameter Lie groups of point transformations with infinitesimal generators in the characteristic form $\mathbf{v} = Q(x, t, u, u_x, u_t) \partial_u$ to one-parameter s -order local transformations with infinitesimal generators of the form

$$\mathbf{v} = Q(x, t, u, \partial u, \partial^2 u, \dots, \partial^s u) \partial_u, \tag{20}$$

where the infinitesimal components depend on derivatives of u up to some finite order $s \geq 1$. The prolongation of \mathbf{v} is given by

$$\mathbf{v}^{(\infty)} = Q \partial_u + D_x Q \partial_{u_x} + D_t Q \partial_{u_t} + D_x^2 Q \partial_{u_{xx}} + D_x D_t Q \partial_{u_{xt}} + D_t^2 Q \partial_{u_{tt}} + \dots \tag{21}$$

where D_x and D_t are total derivative w.r.t x and t , respectively [1].

Then, for $s = 3$, (21) is an infinitesimal local symmetry of the (1) if and only if

$$\begin{aligned} \mathbf{v}^{(\infty)} \left(u_{xt} - \alpha u - \frac{1}{3} \beta (u^3)_{xx} \right) &= 0, & u_{xt} &= \alpha u + \frac{1}{3} \beta (u^3)_{xx}, \\ u_{x^2 t} = D_x \left(\alpha u + \frac{1}{3} \beta (u^3)_{xx} \right), & & u_{xt^2} &= D_t \left(\alpha u + \frac{1}{3} \beta (u^3)_{xx} \right), \\ \dots \dots \dots & & u_{xtt} &= D_t^2 \left(\alpha u + \frac{1}{3} \beta (u^3)_{xx} \right), \end{aligned} \tag{22}$$

which leads to a polynomial of u_{x^5} and u_{t^5} , with functional coefficients of

$$Q(x, t, u, u_x, u_t, u_{xx}, u_{tt}, u_{xxx}, u_{ttt}, u_{x^4}, u_{t^4}) \tag{23}$$

and its derivatives. All of its coefficients must be zero. This leads to a system of 5 linear determining PDEs:

$$\begin{aligned} \beta^4 u_{xx}^8 Q_{u_{t^4}} + Q_{u_{x^4}, u_{t^4}} &= 0, \\ \dots \dots \dots & \\ u_x u_{tt} Q_{u, u_t} + \dots + u_x u_{t^4} Q_{u, u_{ttt}} &= 0. \end{aligned} \tag{24}$$

Therefore, the most general third-order characteristic function Q is

$$\begin{aligned} Q = (c_1 t + c_2) u_t + 3c_1 u - c_1 x u_x + c_3 u_{ttt} - c_3 \beta^3 u_{xx}^6 u_{xxx} & \\ - \frac{3}{2} c_3 \alpha \beta^2 u_x u_{xx}^4 - (c_3 \beta \alpha^2 u_x^2 - c_5) u_x + \frac{c_4 u_{xxx}}{\sqrt{2\beta u_{xx}^2 + \alpha}}, & \end{aligned} \tag{25}$$

where c_1, \dots, c_5 are arbitrary constants. There is not any non-trivial second or fourth-order characteristics. Thus, we prove that

Theorem 3. *The most general third-order infinitesimal local symmetry generator of SPE (1) is a \mathbb{R} -linear combination of following five vector fields $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ of (10) and*

$$\begin{aligned} \mathbf{v}_4 &= \frac{u_{xxx}}{\sqrt{2\beta u_{xx}^2 + \alpha}} \partial_u, \\ \mathbf{v}_5 &= \left(u_{xxx} - \beta^3 u_{xx}^6 u_{xxx} - \frac{3}{2} \alpha \beta^2 u_x u_{xx}^4 - \alpha^2 \beta u_{xxx} \right) \partial_u. \end{aligned} \tag{26}$$

There is not any non-trivial second or fourth-order infinitesimal local symmetry generators.

References

[1] G. W. Bluman, A. F. Cheviakov, S. C. Anco, *Applications of Symmetry Methods to Partial Differential Equations*, Springer, New York, 2010.
 [2] H. Liu, J. Li, Lie symmetry analysis and exact solutions for the short pulse equation, *Nonlinear Analysis* 71 (2009) 2126-2133.
 [3] P. J. Olver, *Applications of Lie Groups to Differential equations*, Second Edition, GTM, Vol. 107, Springer Verlage, New York, 1993.
 [4] P. J. Olver, *Equivalence, Invariant and Symmetry*, Cambridge University Press, Cambridge University Press, Cambridge 1995.
 [5] L.V. Ovsiannikov, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.
 [6] E. Parkes, Some periodic and solitary travelling-wave solutions of the short pulse equation, *Chaos, Solitons & Fractals*, 38 (2008) 154-159.

- [7] A. Sakovich, S. Sakovich, Solitary wave solutions of the short pulse equation, *J. Phys. A Math. Gen.* 39 (2006) L361-L367.
- [8] T. Schafer, C. E. Wayne, Propagation of ultra-short optical pulses in cubic nonlinear media, *Phys. D* 196 (2004) 90-105.

Please cite this article using:

Mehdi Nadjafikhah, Lie group analysis for short pulse equation, *AUT J. Math. Com.*, 1(2) (2020) 223-227
DOI: 10.22060/ajmc.2020.18416.1032

