



Original Article

On bijectivity and Connes bijectivity of a dual Banach algebra with respect to a w^* -closed ideal

Amir Sahami^{*a}, S. Fatemeh Shariati^b, Mehdi Rostami^b, Mona Aj^c

^aDepartment of Mathematics Faculty of Basic Sciences, Ilam University, P.O. Box 69315-516 Ilam, Iran

^bFaculty of Mathematics and Computer Science, Amirkabir University of Technology (Tehran Polytechnic), Iran

^cDepartment of Mathematics, Farhangian University of Kermanshah, Kermanshah, Iran

ABSTRACT: In this paper, we introduce a notion of Connes bijectivity for a dual Banach algebra A with respect to its w^* -closed ideal I , say I -Connes bijectivity. Some Lipschitz algebras $Lip_\alpha(X)$ and some matrix algebras are studied under this new notion. Also, with some mild assumptions, the relation between I -Connes bijectivity and left ϕ -contractibility is given, where ϕ is a w^* -continuous multiplicative linear functional on A . As an application, we characterize Connes bijectivity of some Lipschitz algebras.

Review History:

Received:22 March 2023

Revised:12 May2023

Accepted:16 June 2023

Available Online:01 July 2024

Keywords:

I -Connes bijectivity

left ϕ -contractibility

Dual Banach algebras

MSC (2020):

46M10; 46H25; 46A03

1. Introduction and Preliminaries

The concept of amenability for Banach algebras were first introduced by B. E. Johnson [5]. A Banach algebra A is amenable if and only if there exists a bounded net (m_α) in $A \otimes_p A$ such that $a \cdot m_\alpha - m_\alpha \cdot a \rightarrow 0$ and $\pi_A(m_\alpha)a \rightarrow a$ for every $a \in A$, where $\pi_A : A \otimes_p A \rightarrow A$ is denoted for the product morphism($\pi_A(a \otimes b) = ab$, for all $a, b \in A$). Indeed for a locally compact group G , $L^1(G)$ (the measure algebra $M(G)$) is amenable if and only if G is amenable (G is discrete and amenable). Helemskii in [4] and [15] studied the structure of Banach algebras thorough the homological methods of Banach algebras. He defined the concepts biflatness and bijectivity. In fact a Banach algebra A is bijective, if there exists a bounded A -bimodule morphism $\rho : A \rightarrow A \otimes_p A$ such that $\pi_A \circ \rho(a) = a$, for all $a \in A$. It is known that for a locally compact group G , the group algebra $L^1(G)$ (the measure algebra $M(G)$) is bijective if and only if G is compact (G is finite). For the history of amenability and homological properties of algebra, see [12].

There exists a class of Banach algebras which is called dual Banach algebras. This category of Banach algebras is defined by Runde [11]. Let A be a Banach algebra. Then a Banach A -bimodule E is called dual if there is a

*Corresponding author.

E-mail addresses: a.sahami@ilam.ac.ir, f.shariati@aut.ac.ir, mross@aut.ac.ir, monaaaj1373@yahoo.com



closed submodule E_* of E^* such that $E = (E_*)^*$. The Banach algebra A is called dual if it is dual as a Banach A -bimodule. A dual Banach A -bimodule E is normal if for each $x \in E$ the module maps $A \rightarrow E$ by $a \mapsto a \cdot x$ and $a \mapsto x \cdot a$ are w^* - w^* -continuous. Let A be a Banach algebra and let E be a Banach A -bimodule. A bounded linear map $D : A \rightarrow E$ is called a bounded derivation if $D(ab) = a \cdot D(b) + D(a) \cdot b$, for every $a, b \in A$. A bounded derivation $D : A \rightarrow E$ is called inner if there exists an element x in E such that $D(a) = a \cdot x - x \cdot a$ ($a \in A$). A dual Banach algebra A is called Connes amenable if for every normal dual Banach A -bimodule E , every w^* -continuous derivation $D : A \rightarrow E$ is inner. For a given dual Banach algebra A and a Banach A -bimodule E , $\sigma wc(E)$ denote the set of all elements $x \in E$ such that the module maps $A \rightarrow E$ by $a \mapsto a \cdot x$ and $a \mapsto x \cdot a$ are w^* - w -continuous. It is a closed submodule of E , see [11] and [13] for more details. Note that, since $\sigma wc(A_*) = A_*$, the adjoint of π_A maps A_* into $\sigma wc(A \otimes_p A)^*$. Therefore π_A^{**} drops to an A -bimodule morphism $\pi_{\sigma wc} : (\sigma wc(A \otimes_p A))^* \rightarrow A$.

A dual Banach algebra A is called Connes-biprojective if there exists a bounded A -bimodule morphism $\rho : A \rightarrow (\sigma wc(A \otimes_p A))^*$ such that $\pi_{\sigma wc} \circ \rho(a) = a$ for all $a \in A$. Shirinkalam and Pourabbas showed that a dual Banach algebra A is Connes amenable if and only if A is Connes-biprojective and it has an identity [16]. They characterized Connes-biprojectivity of the measure algebra $M(G)$ for a locally compact group G .

In this paper, we introduce the notion of I -Connes biprojective Banach algebras in the category of dual Banach algebras, where I is a w^* -closed ideal. Some matrix algebras and Lipschitz algebras are studied under this new notion. Also, with some mild assumptions, the relation between I -Connes biprojectivity and left ϕ -contractibility is given, where ϕ is a w^* -continuous multiplicative linear functional on A . As an application, we characterize Connes biprojectivity of some Lipschitz algebras.

Recently the notion of I -biprojectivity is given for Banach algebras. Let A be a Banach algebra and I be a closed ideal of A . Then A is called I -biprojective if there exists a bounded A -bimodule morphism $\rho : I \rightarrow A \otimes_p A$ such that $\pi_A \circ \rho(i) = i$ for all $i \in I$. For a locally compact group G , the measure algebra $M(G)$ is $L^1(G)$ -biprojective if and only if G is compact [14].

Throughout this paper, $\Delta(A)$ ($\Delta_{w^*}(A)$) denotes the character space (w^* - character space) of A , that is, all non-zero (w^* - continuous) multiplicative linear functionals on A , respectively. Let $\phi \in \Delta(A)$. Then ϕ has a unique extension to A^{**} denoted by $\hat{\phi}$ and defined by $\hat{\phi}(F) = F(\phi)$ for every $F \in A^{**}$. Clearly, this extension remains to be a character on A^{**} . The projective tensor product $A \otimes_p A$ is a Banach A -bimodule by the following actions

$$a \cdot (b \otimes c) = ab \otimes c, \quad (b \otimes c) \cdot a = b \otimes ca, \quad (a, b, c \in A).$$

Let X and Y be Banach A -bimodules. Then the map $T : X \rightarrow Y$ is called A -bimodule morphism if

$$T(a \cdot x) = a \cdot T(x), \quad T(x \cdot a) = T(x) \cdot a, \quad (a \in A, x \in X).$$

2. I -Connes biprojectivity

We commence this section with the definition of our new notion. We should remind that every w^* -closed ideal of a dual Banach algebra is also dual Banach algebra see [8, Lemma 2].

Definition 2.1. Let I be a w^* -closed ideal of a dual Banach algebra A . We say that A is I -Connes biprojective, if there exists a bounded A -bimodule morphisms $\rho : I \rightarrow (\sigma wc((A \otimes_p A)^*))^*$ such that $\pi_{\sigma wc} \circ \rho(i) = i$ for all $i \in I$. We say that A is ideally Connes biprojective, if it is I -Connes biprojective for every w^* -closed ideal I of A .

Remark 2.2. In above definition, we can replace ρ with a bounded net of A -bimodule morphisms, say (ρ_α) from I into $\sigma wc((A \otimes_p A)^*)^*$ which $\pi_{\sigma wc} \circ \rho_\alpha(i) \xrightarrow{w^*} i$, for all $i \in I$. To see this, since (ρ_α) is a bounded net of A -bimodule morphisms, we have $(\rho_\alpha) \subseteq B(I, (\sigma wc((A \otimes_p A)^*))^*)$ (the set of bounded linear maps from I into $(\sigma wc((A \otimes_p A)^*))^*$). On the other hand, on bounded sets the w^* -operator topology coincides with the w^* -topology of $B(I, (\sigma wc((A \otimes_p A)^*))^*)$ where identified with $(I \otimes_p \sigma wc(A \otimes_p A))^*$. It is known that the unit ball of $B(I, (\sigma wc((A \otimes_p A)^*))^*)$ is w^* -operator compact. Then (ρ_α) has a w^* -operator topology limit point say ρ . Thus

$$\rho(i_1 i_2) = w^* - \lim \rho_\alpha(i_1 i_2) = w^* - \lim i_1 \cdot \rho_\alpha(i_2) = i_1 \cdot w^* - \lim \rho_\alpha(i_2) = i_1 \cdot \rho(i_2).$$

Similarly we have $\rho(i_1 i_2) = \rho(i_1) \cdot i_2$. Therefore ρ is a bounded A -bimodule morphism and

$$\pi_{\sigma wc} \circ \rho(i) = \pi_{\sigma wc}(w^* - \lim \rho_\alpha(i)) = w^* - \lim \pi_{\sigma wc} \rho_\alpha(i) = i.$$

So A is I -Connes biprojective.

Lemma 2.3. Let A be a non-zero dual Banach algebra and $ab = ba = 0$ for all $a, b \in A$. Then A is not Connes biprojective.

Proof. We assume in contradiction that A is Connes biprojective. Then there exists a bounded A -bimodule morphisms $\rho : A \rightarrow (\sigma wc(A \otimes_p A))^*$ such that $\pi_{\sigma wc} \circ \rho(a) = a$ for each $a \in A$. It is known that there exists a net (u_α) in $A \otimes_p A$ such that $w^* - \lim \hat{u}_\alpha|_{\sigma wc(A \otimes_p A)^*} = \rho(a)$. Thus

$$a = \pi_{\sigma wc} \circ \rho(a) = w^* - \lim \pi_A(u_\alpha) = w^* - \lim 0 = 0.$$

It follows that $A = 0$ which is a contradiction. □

Let A be a Banach algebra and $\phi \in \Delta(A)$. Then A is called left ϕ -contractible if there exists a $m \in A$ such that $am = \phi(a)m$ and $\phi(m) = 1$ for all $a \in A$. For further information about this concept see [10].

Proposition 2.4. *Let A be a commutative dual Banach algebra and $\phi \in \Delta_{w^*}(A)$. Suppose that I is a w^* -closed ideal of A . If A is I -Connes biprojective, then A is left ϕ -contractible, provided that $\phi|_I \neq 0$.*

Proof. Let i_0 be an element of I such that $\phi(i_0) = 1$. Since A is I -Connes biprojective, there exists a bounded A -bimodule morphism $\rho : I \rightarrow (\sigma wc((A \otimes_p A)^*))^*$ such that $\pi_{\sigma wc} \circ \rho(i) = i$ for all $i \in I$. Put $m = \rho(i_0)$. One can see that $a \cdot m = m \cdot a$ and $\pi_{\sigma wc}(m)i = i$ for all $i \in I$ and $a \in A$. Define $T : A \otimes_p A \rightarrow A$ by $T(a \otimes b) = \phi(b)a$ for each $a, b \in A$. It is easy to see that

$$aT(x) = T(a \cdot x), \quad T(x \cdot a) = \phi(a)T(x) \quad \phi \circ T(x) = \phi \circ \pi_A(x) \quad (a \in A, x \in X).$$

Since T^{**} is a w^* -continuous map, for each $x \in (A \otimes_p A)^{**}$ and $a \in A$, we have

$$aT^{**}(x) = T^{**}(a \cdot x), \quad T^{**}(x \cdot a) = \phi(a)T^{**}(x) \quad \tilde{\phi} \circ T^{**}(x) = \tilde{\phi} \circ \pi_A^{**}(x).$$

On the other hand, for all $a \in A$, $f \in A^*$ and $x \in A \otimes_p A$ consider

$$\langle x, a \cdot T^*(f) \rangle = \langle x \cdot a, T^*(f) \rangle = \langle T(x \cdot a), f \rangle = \phi(a)\langle T(x), f \rangle = \phi(a)\langle x, T^*(f) \rangle$$

also

$$\langle x, T^*(f) \cdot a \rangle = \langle a \cdot x, T^*(f) \rangle = \langle T(a \cdot x), f \rangle = \langle a \cdot T(x), f \rangle = \langle T(x), f \cdot a \rangle,$$

which follow that

$$a \cdot T^*(f) = \phi(a)T^*(f), \quad T^*(f) \cdot a = T^*(f \cdot a).$$

These last facts with the w^* -continuity of ϕ gives that $T^*(\sigma wc(A^*)) \subseteq \sigma wc(A \otimes_p A)^*$. Let $q : A^{**} \rightarrow \sigma wc((A)^*)^*$ be the quotient map. For each $a \in A$ and $f \in \sigma wc(A^*)$, we have

$$\begin{aligned} \langle f \cdot a, q \circ T^{**}(m) \rangle &= \langle f \cdot a, T^{**}(m)|_{\sigma wc(A)^*} \rangle = \langle f \cdot a, T^{**}(m) \rangle \\ &= \langle f, a \cdot T^{**}(m) \rangle \\ &= \langle f, T^{**}(a \cdot m) \rangle \\ &= \langle T^*(f), a \cdot m \rangle \\ &= \langle T^*(f), m \cdot a \rangle \\ &= \langle f, T^{**}(m \cdot a) \rangle \\ &= \langle f, \phi(a)T^{**}(m) \rangle \\ &= \phi(a)\langle f, T^{**}(m) \rangle \\ &= \phi(a)\langle f, T^{**}(m)|_{\sigma wc(A)^*} \rangle \\ &= \phi(a)\langle f, q \circ T^{**}(m) \rangle. \end{aligned}$$

It follows that $a \cdot q \circ T^{**}(m) = \phi(a)q \circ T^{**}(m)$ for all $a \in A$. Moreover we know that ϕ is a w^* -multiplicative linear functional. It follows that $\phi \in A_*$. On the other hand $A_* \subseteq \sigma wc(A)^*$. Then $\phi \in \sigma wc(A)^*$. Hence

$$\langle \phi, q \circ T^{**}(m) \rangle = \langle \phi, T^{**}(m)|_{\sigma wc(A)^*} \rangle = \langle \phi, T^{**}(m) \rangle = \langle \phi, \pi_A^{**}(m) \rangle = 1.$$

Thus it gives that A is left ϕ -contractible. □

Theorem 2.5. *Let A be a dual Banach algebra and $\phi \in \Delta_{w^*}(A)$. Suppose that I is a w^* -closed ideal which $\phi|_I \neq 0$ and $I \ker \phi|_I = \ker \phi|_I$. If A is I -Connes biprojective, then A is left ϕ -contractible.*

Proof. Let A be I -Connes biprojective. Then there exists a bounded A -bimodule morphism $\rho : I \rightarrow (\sigma wc(A \otimes_p A))^*$ such that $\pi_{\sigma wc} \circ \rho(i) = i$ for all $i \in I$. Let i_0 be an element of I such that $\phi(i_0) = 1$. It is known that $\ker \phi$ is a closed ideal of A . Thus $\frac{A}{\ker \phi}$ is a Banach A -bimodule, naturally. We denote the identity map on A by id_A . Also $q : A \rightarrow \frac{A}{\ker \phi}$ is denoted for the quotient map. Define $id_A \otimes q : A \otimes_p A \rightarrow A \otimes_p \frac{A}{\ker \phi}$ by $id_A \otimes q(a \otimes b) = a \otimes (b + \ker \phi)$ for all $a, b \in A$. Clearly $id_A \otimes q$ is a bounded A -bimodule morphism. It implies that

$$(id_A \otimes q)^*(\sigma wc(A \otimes_p \frac{A}{\ker \phi}))^* \subseteq \sigma wc(A \otimes_p A)^*.$$

Using this fact, set

$$\theta : ((id_A \otimes q)|_{\sigma wc(A \otimes_p \frac{A}{\ker \phi})})^* : (\sigma wc(A \otimes_p A))^* \rightarrow (\sigma wc(A \otimes_p \frac{A}{\ker \phi}))^*.$$

Clearly we observe that θ is a w^* -continuous A -bimodule morphism. Put

$$\eta = \theta \circ \rho : I \rightarrow (\sigma wc(A \otimes_p \frac{A}{\ker \phi}))^*.$$

We can see that η is a bounded A -bimodule morphism. Since $\overline{I \ker \phi|_I} = I$, we may assume that for each $l \in \ker \phi|_I$ there is $l_1 \in \ker \phi|_I$ and $i_1 \in I$ such that $l = i_1 l_1$. On the other hand we know that there exists a quotient map q from $(A \otimes_p A)^{**} \rightarrow (\sigma wc(A \otimes_p A))^*$ and compose q with the embedding map from $A \otimes_p A$ into $(A \otimes_p A)^{**}$ gives a continuous A -bimodule map $\tau : A \otimes_p A \rightarrow (\sigma wc(A \otimes_p A))^*$ which has a w^* -dense range. We denote \bar{u} for $\tau(u) = \hat{u}|_{\sigma wc(A \otimes_p A)^*}$, where $u \in A \otimes_p A$ and \hat{u} is the image of embedding map at u in $(A \otimes_p A)^{**}$. So for $\rho(i_1) \in (\sigma wc(A \otimes_p A))^*$ there exists a net (u_α) in $A \otimes_p A$ which $w^* - \lim \bar{u}_\alpha = \rho(i_1)$. Applying the w^* -continuity of θ implies that

$$\begin{aligned} \eta(l) &= \theta \circ \rho(i_1 l_1) = \theta(\rho(i_1) \cdot l_1) \\ &= \theta((w^* - \lim \bar{u}_\alpha) \cdot l_1) \\ &= w^* - \lim \theta(\bar{u}_\alpha \cdot l_1) \\ &= w^* - \lim ((id_A \otimes q)|_{\sigma wc(A \otimes_p \frac{A}{\ker \phi})})^*(u_\alpha \cdot l_1) = 0, \end{aligned}$$

the last equality holds because $q(l_1) = 0$. So $\eta(l) = 0$. So η induces a map from $\frac{A}{\ker \phi}$ into $(\sigma wc(A \otimes_p A))^*$ which is a bounded A -bimodule morphism. Since $\phi \in \Delta_{w^*}(A)$, we denote $\bar{\phi} : \frac{A}{\ker \phi} \rightarrow \mathbb{C}$ for a character which is given by $\bar{\phi}(a + \ker \phi) = \phi(a)$ for all $a \in A$. Clearly $\bar{\phi}$ is a character. Put $id_A \otimes \bar{\phi} : A \otimes_p \frac{A}{\ker \phi} \rightarrow A$ which is defined by $id_A \otimes \bar{\phi}(a \otimes b + \ker \phi) = \phi(b)a$ for every $a, b \in A$. One can readily see that for each $f \in A^*$ and $a \in A$

$$(id_A \otimes \bar{\phi})^*(f) \cdot a = (id_A \otimes \bar{\phi})^*(f \cdot a), \quad a \cdot ((id_A \otimes \bar{\phi})^*(f)) = \phi(a)(id_A \otimes \bar{\phi})^*(f).$$

Using w^* -continuity of ϕ and $\sigma wc(A_*) = A_*$ implies that

$$(id_A \otimes \bar{\phi})(A_*) = (id_A \otimes \bar{\phi})(\sigma wc(A_*)) \subseteq \sigma wc(A \otimes_p \frac{A}{\ker \phi})^*.$$

It follows that

$$\psi = ((id_A \otimes \bar{\phi})|_{A_*})^* : (\sigma wc(A \otimes_p \frac{A}{\ker \phi}))^* \rightarrow A$$

is a w^* -continuous left A -module morphism. Set $y = \psi \circ \eta$. Hence y is a bounded left A -module morphism from $\frac{A}{\ker \phi}$ into A . Note that y is a non-zero map. To see this, we show that $\phi \circ \psi = \phi \circ \pi_{\sigma wc}$. Clearly for each $a, b \in A$, we have

$$\begin{aligned} \phi \circ (id_A \otimes \bar{\phi}) \circ (id_A \otimes q)(a \otimes b) &= \phi \circ (id_A \otimes \bar{\phi})(a \otimes (b + \ker \phi)) = \phi(a\phi(b)) \\ &= \phi(a)\phi(b) \\ &= \phi \circ \pi_A. \end{aligned}$$

On the other hand for each $v \in A \otimes_p \frac{A}{\ker \phi}$ we have $\psi(\hat{v}|_{(A \otimes_p \frac{A}{\ker \phi})^*}) = (id_A \otimes \bar{\phi})(v)$. Also, for each $u \in A \otimes_p A$, we have $\pi_{\sigma wc}(\bar{u}) = \pi_A(u)$. Let $m \in (\sigma wc(A \otimes_p A))^*$. Then there exists a net (u_α) in $A \otimes_p A$ such that $m = w^* - \lim \bar{u}_\alpha$.

As we know that ϕ, θ, ψ and $\pi_{\sigma_{wc}}$ are w^* -continuous maps. So

$$\begin{aligned} \phi \circ \psi \circ \theta(m) &= \phi \circ \psi \circ \theta(w^* - \lim \bar{u}_\alpha) = w^* - \lim \phi \circ \psi \circ \theta(\bar{u}_\alpha) \\ &= w^* - \lim \phi \circ (id_A \otimes \bar{\phi}) \circ (id_A \otimes q)(u_\alpha) \\ &= w^* - \lim \phi \circ \pi_A(u_\alpha) \\ &= w^* - \lim \phi \circ \pi_{\sigma_{wc}}(\bar{u}_\alpha) \\ &= \phi \circ \pi_{\sigma_{wc}}(m). \end{aligned}$$

Thus for $i_0 \in I \subseteq A$, we have

$$\begin{aligned} \phi \circ y(i_0 + \ker \phi) &= \phi \circ \psi \circ \eta(i_0 + \ker \phi) = \phi \circ \psi \circ \theta \circ \rho(i_0) \\ &= \phi \circ \pi_{\sigma_{wc}} \circ \rho(i_0) = \phi(i_0) = 1. \end{aligned}$$

It implies that y is nonzero map as desired. Also for each $a \in A$, we have

$$\begin{aligned} ay(i_0 + \ker \phi) &= y(ai_0 + \ker \phi) = y(\phi(a)i_0 + \ker \phi) \\ &= \phi(a)y(i_0 + \ker \phi). \end{aligned}$$

Hence A is left ϕ -contractible. □

We give a dual Banach algebra A with a w^* -closed ideal I which neither A nor I is Connes biprojective. But A is I -Connes biprojective.

Example 2.1. Let $A = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid a, b, c \in \mathbb{C} \right\}$ and $I = \left\{ \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix} \mid b \in \mathbb{C} \right\}$. With matrix operations and the ℓ^1 -norm, A becomes a dual Banach algebra and I becomes a w^* -closed ideal of A . We assume in contradiction that A is Connes biprojective. Since A is unital by [16, Theorem 2.2] A is Connes amenable. Define $\phi : A \rightarrow \mathbb{C}$ by $\phi\left(\begin{pmatrix} 0 & b \\ 0 & c \end{pmatrix}\right) = c$. Clearly ϕ is a character on A . Put $J = \left\{ \begin{pmatrix} 0 & b \\ 0 & c \end{pmatrix} \mid b, c \in \mathbb{C} \right\}$. It is easy to verify that J is a w^* -closed ideal of A which $\phi|_J \neq 0$. It is easy to see that Connes amenability of A implies that A is left ϕ -amenable (or A is left ϕ -contractible), see [7]. So by similar method as in [6, Lemma 3.1] we have J is left ϕ -contractible. That is there is an element $m = \begin{pmatrix} 0 & b_0 \\ 0 & c_0 \end{pmatrix}$ in J such that $jm = \phi(j)m$ and $\phi(m) = 1$ for each $j \in J$, where $b_0, c_0 \in \mathbb{C}$. Suppose that $j = \begin{pmatrix} 0 & j_1 \\ 0 & j_2 \end{pmatrix}$ is an arbitrary element of J , where j_1 and j_2 in \mathbb{C} . Thus

$$jm = \begin{pmatrix} 0 & j_1 \\ 0 & j_2 \end{pmatrix} \begin{pmatrix} 0 & b_0 \\ 0 & c_0 \end{pmatrix} = \begin{pmatrix} 0 & j_1 c_0 \\ 0 & j_2 c_0 \end{pmatrix} = \phi(j)m = j_2 \begin{pmatrix} 0 & b_0 \\ 0 & c_0 \end{pmatrix} = \begin{pmatrix} 0 & j_2 b_0 \\ 0 & j_2 c_0 \end{pmatrix}$$

and $\phi\left(\begin{pmatrix} 0 & b_0 \\ 0 & c_0 \end{pmatrix}\right) = c_0 = 1$. It follows that for each j_1 and j_2 in \mathbb{C} we have $j_2 b_0 = j_1$. Put $j_2 = 0$ and $j_1 = 1$. Then contradiction reveals.

Since for each element i_1 and i_2 in I , we have $i_1 i_2 = 0$, Lemma 2.3 follows that I is not Connes biprojective. To show that A is I -Connes biprojective define $\rho : I \rightarrow A \otimes_p A \subseteq (\sigma_{wc}(A \otimes_p A))^*$ by

$$\rho\left(\begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix}\right) = \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}, \quad (b \in \mathbb{C})$$

Clearly ρ is a bounded A -bimodule morphism and $\pi_{\sigma_{wc}} \circ \rho(i) = i$ for all $i \in I$.

3. Applications for Lipschitz algebras

Let X be a compact metric space and $\alpha > 0$. The space of complex valued function on X is denoted by $Lip_\alpha(X)$ which

$$p_\alpha(f) = \sup\left\{ \frac{|f(x) - f(y)|}{d(x, y)^\alpha} : x, y \in X, x \neq y \right\}$$

is finite. Also

$$lip_\alpha(X) = \{f \in Lip_\alpha(X) : \frac{|f(x) - f(y)|}{d(x, y)^\alpha} \rightarrow 0 \text{ as } d(x, y) \rightarrow 0\}.$$

Define

$$\|f\|_\alpha = \|f\|_\infty + p_\alpha(f).$$

With $\|\cdot\|_\alpha$ and the pointwise operations $Lip_\alpha(X)$ and $lip_\alpha(X)$ become Banach algebras. It is known that for $0 < \alpha < 1$, $lip_\alpha(X)^{**}$ is isometrically isomorphism with $Lip_\alpha(X)$. Also $Lip_\alpha(X)$ and $lip_\alpha(X)$ are Arens regular Banach algebras for more details, see [2]. Recently Minapour and Zivari-Kazmpour showed that $Lip_\alpha(X)$ is a dual Banach algebra, [9].

Theorem 3.1. *Let X be a compact metric space and $0 < \alpha < 1$. Then $Lip_\alpha(X)$ is Connes biprojective if and only if X is finite.*

Proof. Suppose that $Lip_\alpha(X)$ is Connes biprojective. Since $Lip_\alpha(X)$ posses an identity, by [16, Theorem 2.2] $Lip_\alpha(X)$ is Connes amenable. Thus $Lip_\alpha(X) \cong (lip_\alpha(X))^{**}$ is Connes amenable. It is easy to see that $lip_\alpha(X)$ is a closed ideal of $Lip_\alpha(X)$. Applying [12, Theorem 4.4.8] follows that $lip_\alpha(X)$ is amenable. By the main result of [3] X is finite.

Converse is clear. □

A Banach algebra A is called biflat if there exists a bounded A -bimodule morphism $\rho : A \rightarrow (A \otimes_p A)^{**}$ such that $\pi_A^{**} \circ \rho(a) = a$ for each $a \in A$ [12].

Theorem 3.2. *Let X be a compact metric space and $0 < \alpha < 1$. Then $Lip_\alpha(X)$ is $lip_\alpha(X)$ -biprojective if and only if X is finite.*

Proof. Suppose that $Lip_\alpha(X)$ is $lip_\alpha(X)$ -biprojective. Then by [14, Lemma 3.5] $lip_\alpha(X)$ is biflat. Clearly $lip_\alpha(X)$ posses an identity. Thus $lip_\alpha(X)$ is amenable. Applying [3] X is finite.

For converse, let X be finite. Then by [1, Corollary 2.2] $Lip_\alpha(X)$ separates the point of X . Applying [1, Theorem 3.2] follows that $Lip_\alpha(X)$ is biprojective. Then there exists a bounded $Lip_\alpha(X)$ -bimodule morphism ρ from $Lip_\alpha(X)$ into $Lip_\alpha(X) \otimes_p Lip_\alpha(X)$ such that $\pi_{Lip_\alpha(X)} \circ \rho(a) = a$ for all $a \in Lip_\alpha(X)$. Restrict ρ on $lip_\alpha(X)$ finishes the proof. □

Let X be a metric space. A subalgebra A of $C_b(X)$ (Banach algebra of bounded and continuous functions) is called strongly separating the points of X , if for each $x, y \in X$ with $x \neq y$, there exists $f \in A$ such that $f(x) \neq f(y)$.

Proposition 3.3. *Let G be a metric space which is a compact group and $\alpha > 0$. Suppose that $Lip_\alpha(G)$ is strongly separating the points of G . Let I be a non-zero closed ideal of $Lip_\alpha(G)$. Then $Lip_\alpha(G)$ is I -biprojective if and only if G finite.*

Proof. Since I is a non-zero closed ideal of $Lip_\alpha(G)$, semisimplicity of $Lip_\alpha(G)$ gives that there exists a non-zero multiplicative linear functional ϕ_g on $Lip_\alpha(G)$ such that $\phi_g|_I \neq 0$. By some modifications of the arguments as in Proposition 2.4, I -biprojectivity of $Lip_\alpha(G)$ implies that $Lip_\alpha(G)$ is left ϕ_g -contractible. So there exists $m \in Lip_\alpha(G)$ such that $fm = \phi_g(f)m$ and $\phi_g(m) = m(g) = 1$ for all $f \in Lip_\alpha(G)$. Let $y \neq g$ be an arbitrary element of G . Since $Lip_\alpha(G)$ is strongly separating the element of G , by [1, Proposition 2.1] there exists a $f_0 \in Lip_\alpha(G)$ such that $f_0(g) = 1$ and $f_0(y) = 0$. we know that $m = \phi_g(f_0)m = f_0m$. Thus $m(y) = f_0m(y) = f_0(y)m(y) = 0$ and $m(g) = f_0m(g) = f_0(g)m(g) = 1$. So $m = \chi_{\{g\}} \in Lip_\alpha(G)$, the characteristic function at g . Since m is a continuous function on G , it gives that G is discrete (and compact). Then G is finite.

The converse is similar to the only if part of previous Theorem. □

Acknowledgement

The authors would like to thank the anonymous referees for their very careful reading and their valuable comments which improved the manuscript. Also the first author thanks Ilam university for its support.

References

- [1] F. ABTAHI, E. BYABANI, AND A. REJALI, *Some algebraic and homological properties of Lipschitz algebras and their second duals*, Arch. Math. (Brno), 55 (2019), pp. 211–224.
- [2] W. G. BADE, P. C. CURTIS, JR., AND H. G. DALES, *Amenability and weak amenability for Beurling and Lipschitz algebras*, Proc. London Math. Soc. (3), 55 (1987), pp. 359–377.
- [3] F. GOURDEAU, *Amenability of Banach algebras*, Math. Proc. Cambridge Philos. Soc., 105 (1989), pp. 351–355.

- [4] A. Y. HELEMSKII, *The homology of Banach and topological algebras*, vol. 41 of Mathematics and its Applications (Soviet Series), Kluwer Academic Publishers Group, Dordrecht, 1989. Translated from the Russian by Alan West.
- [5] B. E. JOHNSON, *Cohomology in Banach algebras*, Memoirs of the American Mathematical Society, No. 127, American Mathematical Society, Providence, RI, 1972.
- [6] E. KANIUTH, A. T. LAU, AND J. PYM, *On ϕ -amenability of Banach algebras*, Math. Proc. Cambridge Philos. Soc., 144 (2008), pp. 85–96.
- [7] A. MAHMOODI, *φ -contractibility and φ -Connes amenability coincide with some older notions*, Bull. Aust. Math. Soc., 97 (2018), pp. 274–278.
- [8] A. MINAPOOR, A. BODAGHI, AND D. EBRAHIMI BAGHA, *Ideal Connes-amenability of dual Banach algebras*, Mediterr. J. Math., 14 (2017), pp. Paper No. 174, 12.
- [9] A. MINAPOOR AND A. ZIVARI-KAZEMPOUR, *Ideal Connes-amenability of certain dual Banach algebras*, Complex Anal. Oper. Theory, 17 (2023), pp. Paper No. 27, 11.
- [10] R. NASR-ISFAHANI AND S. SOLTANI RENANI, *Character contractibility of Banach algebras and homological properties of Banach modules*, Studia Math., 202 (2011), pp. 205–225.
- [11] V. RUNDE, *Amenability for dual Banach algebras*, Studia Math., 148 (2001), pp. 47–66.
- [12] ———, *Lectures on amenability*, vol. 1774 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 2002.
- [13] ———, *Dual Banach algebras: Connes-amenability, normal, virtual diagonals, and injectivity of the predual bimodule*, Math. Scand., 95 (2004), pp. 124–144.
- [14] A. SAHAMI, M. ROSTAMI, AND S. KALANTARI, *On I -biflat and I -biprojective Banach algebras*, Wavelet and Linear Algebra, 8 (2021), pp. 49–59.
- [15] A. SAHLEH AND A. Z. KAZEMPOUR, *Some results on Banach precompact elements of a locally convex algebra*, Far East J. Math. Sci. (FJMS), 31 (2008), pp. 17–23.
- [16] A. SHIRINKALAM AND A. POURABBAS, *Connes-biprojective dual Banach algebras*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys., 78 (2016), pp. 175–184.

Please cite this article using:

Amir Sahami, S. Fatemeh Shariati, Mehdi Rostami, Mona Aj, On biprojectivity and Connes biprojectivity of a dual Banach algebra with respect to a w^* -closed ideal, AUT J. Math. Com., 6(1) (2024) 27-33
<https://doi.org/10.22060/AJMC.2023.22285.1149>

